

The Stueckelberg Z' Extension with Kinetic Mixing and Milli-Charged Dark Matter from the Hidden Sector

Daniel Feldman,^{1,*} Zuowei Liu,^{1,†} and Pran Nath^{1,‡}

¹*Department of Physics, Northeastern University, Boston, MA 02115, USA*

V1: February 12, 2007 ; Published: June 1, 2007, Phys. Rev. D 75, 115001 2007

An analysis is given of the Stueckelberg extension of the Standard Model with a hidden sector gauge group $U(1)_X$ where the mass growth for the extra gauge boson occurs via the Stueckelberg mechanism, and where the kinetic mixing in the $U(1)_X \times U(1)_Y$ sector is included. Such a kinetic mixing is generic in a broad class of supergravity and string models. We carry out a detailed global fit of the model with the precision LEP data on and off the Z pole, with χ^2 within 1% of the χ^2 of the Standard Model fit. Further, it is shown that in the absence of matter in the hidden sector, there is a single effective parameter that controls the deviations from the Standard Model predictions, and the dependence on the kinetic mixing emerges only when matter in the hidden sector is included. An analysis is also given of milli-charged dark matter arising from the hidden sector, where it is shown that such dark matter from the Stueckelberg extension can satisfy WMAP-3 data while allowing for a sharp Z' resonance which can be detected at the Tevatron and at the LHC via a dilepton signal generated by the Drell-Yan process.

I. INTRODUCTION

Recent works on the Stueckelberg extension of the SM [1] (StSM) and of the MSSM [2, 3] (StMSSM), have shown consistency with the LEP data while allowing for the possibility of a narrow resonance which could lie as low as just above the Z mass [1, 2, 3, 4, 5]. Similar phenomena regarding a narrow resonance are seen in other classes of models such as those based on universal extra dimensions [6], models with a shadow sector [7, 8, 9], and in the models considered in Ref. [10]. The possibility of a narrow graviton resonance also arises in the RS model [11, 12]. Thus the study of narrow resonances is a topic of significant interest. In this work we focus on the Stueckelberg extensions further. Such models may be a field theory realization of string models arising from orientifolds [13, 14, 15, 16, 17] and their discovery could be harbingers of a new regime of physics altogether. The supersymmetric Stueckelberg extensions have the possibility of generating a new type of neutral Majorana fermion which is massive and extra-weakly interacting, and with R parity conservation, a candidate for cold dark matter [18]. Indeed a detailed analysis shows that the relic density in such models is consistent with the three year WMAP data [19]. Additionally, the Stueckelberg extensions give rise to milli-charges for matter residing in the hidden sector [1, 3]. A recent analysis [20] indicates that such matter can annihilate in sufficient amounts to satisfy the relic density constraints from WMAP-3 for a broad Z' resonance.

The main focus of this work is an extension of the

class of models considered in Refs. [1, 2, 3, 4, 5] by including kinetic mixing between the two Abelian $U(1)$ gauge fields. Specifically we consider the extended electroweak sector with the gauge groups $SU(2)_L \times U(1)_Y \times U(1)_X$ where the Stueckelberg mechanism along with the spontaneous breaking in the Higgs sector generates the vector boson mass, and a mixing in the gauge kinetic energy of the $U(1)_X \times U(1)_Y$ sector is included. Such kinetic mixings can arise in a variety of ways [21, 22] and enter prominently in models of the type considered in Refs. [8, 9]. The model considered here encompasses the models of Refs. [1, 2, 3, 4, 5] which can be obtained in certain limits of the model discussed here. Inclusion of the kinetic mixing in the Stueckelberg extension enhances significantly the parameter space where new physics can exist consistent with the stringent LEP, Tevatron, and WMAP constraints. This parameter space includes the possibility of a narrow Z' resonance very distinct from the Z' of the conventional models [23, 24, 25].

The outline of the rest of the paper is as follows: In Sec.(II) we give a description of the Stueckelberg extension with kinetic mixing in the $U(1)_X \times U(1)_Y$ sector. In Sec.(III) we give a detailed numerical analysis of the electroweak constraints from LEPI and LEPII. In Sec.(IV) an analysis of milli-charged dark matter that arises from the hidden sector is given. Conclusions are given in Sec.(V). Appendices contain several mathematical details, including an analysis regarding the origin of milli-charged matter.

II. THE STUECKELBERG Z' EXTENSION WITH KINETIC MIXING

In this section we discuss the $U(1)_X$ Stueckelberg [26] extension of the Standard Model (SM) with gauge

*Electronic address: feldman.da@neu.edu

†Electronic address: liu.zu@neu.edu

‡Electronic address: nath@lepton.neu.edu

kinetic mixing (StkSM). We assume that the quarks, leptons and the Higgs field of the SM do not carry $U(1)_X$ quantum numbers, and the fields in the hidden sector do not carry quantum numbers of the SM gauge group. Thus the $U(1)_X$ sector is hidden except for the $U(1)_X \times U(1)_Y$ mixings in the gauge vector boson sector as given by the effective Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{StkSM}} &= \mathcal{L}_{\text{SM}} + \Delta\mathcal{L}, \\ \Delta\mathcal{L} &\supset -\frac{1}{4}C_{\mu\nu}C^{\mu\nu} - \frac{\delta}{2}C_{\mu\nu}B^{\mu\nu} \\ &\quad - \frac{1}{2}(\partial_\mu\sigma + M_1C_\mu + M_2B_\mu)^2 + g_X J_X^\mu C_\mu. \end{aligned} \quad (1)$$

Here B_μ is the $U(1)_Y$ gauge field, C_μ is the $U(1)_X$ gauge field with coupling strength g_X to hidden matter through the source J_X , σ is the axion, M_1 and M_2 are mass parameters, and δ is the kinetic mixing parameter. The σ field is charged under both $U(1)_X$ and $U(1)_Y$ and the Lagrangian of Eq. (1) is invariant under the $U(1)_X \times U(1)_Y$ gauge transformations

$$\begin{aligned} \delta B_\mu &= \partial_\mu\lambda_X, \quad \delta C_\mu = 0, \quad \delta\sigma = -M_2\lambda_X \\ \delta B_\mu &= 0, \quad \delta C_\mu = \partial_\mu\lambda_Y, \quad \delta\sigma = -M_1\lambda_Y. \end{aligned} \quad (2)$$

The above model has a non-diagonal kinetic mixing matrix ($\mathcal{K}$) and a non-diagonal mass matrix (M_{St}^2) and in the unitary gauge [1] in the basis $V^T = (C, B, A^3)$

$$\mathcal{K} = \begin{bmatrix} 1 & \delta & 0 \\ \delta & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad (3)$$

$$M_{\text{St}}^2 = \begin{bmatrix} M_1^2 & M_1M_2 & 0 \\ M_1M_2 & M_2^2 + \frac{1}{4}v^2g_Y^2 & -\frac{1}{4}v^2g_2g_Y \\ 0 & -\frac{1}{4}v^2g_2g_Y & \frac{1}{4}v^2g_2^2 \end{bmatrix}. \quad (4)$$

A simultaneous diagonalization of the kinetic energy and of the mass matrix can be obtained by a transformation $T = KR$, which is a combination of a $GL(3)$ transformation (K) and an orthogonal transformation (R). This allows one to work in the diagonal basis, denoted by E where $E^T = (Z', Z, A_\gamma)$, through the transformation $V = (KR)E$, where the matrix K

which diagonalizes the kinetic terms has the form

$$K = \begin{bmatrix} C_\delta & 0 & 0 \\ -S_\delta & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad C_\delta = \frac{1}{\sqrt{1-\delta^2}}, \quad S_\delta = \delta C_\delta. \quad (5)$$

The matrix R is then defined by the diagonalization of the mass matrix

$$M_D^2 = R^T (K^T M_{\text{St}}^2 K) R. \quad (6)$$

The model of Eq.(1) involves three parameters: M_1, M_2, δ or alternately M_1, ϵ, δ where $\epsilon = M_2/M_1$. Expressing the transformation $T = KR$ in terms of the matrix elements, one has

$$\begin{aligned} T_{1j} &= C_\delta R_{1j}, \\ T_{2j} &= R_{2j} - S_\delta R_{1j}, \\ T_{3j} &= R_{3j}. \end{aligned} \quad (7)$$

The neutral current interaction with the visible sector fermions is given by

$$\mathcal{L}_{NC} = -\frac{1}{i} \sum_f [\bar{f}_L D_\mu \gamma^\mu f_L + (L \rightarrow R)], \quad (8)$$

where D_μ is the covariant derivative with respect to $SU(2)_L \times U(1)_Y \times U(1)_X$ gauge group except that, as mentioned in the beginning, we assume that the visible sector matter, i.e., quarks, leptons and the Higgs, are not charged under $U(1)_X$. Thus the covariant derivative includes only the $SU(2)_L$ gauge coupling g_2 and $U(1)_Y$ gauge coupling g_Y . The diagonalization also leads to the following relation for the electronic charge

$$\frac{1}{e^2} = \frac{1}{g_2^2} + \frac{1 - 2\epsilon\delta + \epsilon^2}{g_Y^2}. \quad (9)$$

Thus g_Y is related to g_Y^{SM} by

$$g_Y = \gamma \sqrt{1 + \epsilon^2 - 2\delta\epsilon}, \quad \gamma \equiv g_Y^{SM}, \quad (10)$$

and one may write the neutral current interaction so that

$$\mathcal{L}_{NC} = \frac{\sqrt{g_2^2 + \gamma^2}}{2} \bar{f} \gamma^\mu [(v'_f - \gamma_5 a'_f) Z'_\mu + (v_f - \gamma_5 a_f) Z_\mu] f + e \bar{f} \gamma^\mu Q_f A_\mu f, \quad (11)$$

where

$$\begin{aligned} v_f &= \frac{1}{\sqrt{g_2^2 + \gamma^2}} [(g_2 T_{32} - g_Y T_{22}) T_f^3 + 2g_Y T_{22} Q_f], \\ a_f &= \frac{1}{\sqrt{g_2^2 + \gamma^2}} [(g_2 T_{32} - g_Y T_{22}) T_f^3], \\ v'_f &= \frac{1}{\sqrt{g_2^2 + \gamma^2}} [(g_2 T_{31} - g_Y T_{21}) T_f^3 + 2g_Y T_{21} Q_f], \\ a'_f &= \frac{1}{\sqrt{g_2^2 + \gamma^2}} [(g_2 T_{31} - g_Y T_{21}) T_f^3], \end{aligned} \quad (12)$$

and where, as usual, $Q_f = T_f^3 + Y_f/2$. We note that the mass and kinetic mixing parameters enter not only through T_{ij} but also through g_Y via the constraint of Eq.(9).




FIG. 1: An analysis of χ^2 in StkSM model in the $\epsilon - \delta$ plane. The center white region is where χ^2 of the StkSM model is within 1% of the SM fits. Along the line $\epsilon = \delta$ the Z' decouples and the model gives the same χ^2 fit to data as in the SM (see also Appendix A).

III. CONSTRAINTS FROM ELECTROWEAK DATA

We discuss now the constraints on the model with both mass mixing and kinetic mixing from the precision electroweak data. We start by assuming that the hidden sector does not contain matter, and the case when matter is included in the hidden sector is discussed in Sec.(IV) and in Appendix B. To obtain the allowed range of ϵ and δ , we follow the same approach as in Ref. [4, 5, 27]. The first constraint comes from the comparison of the one sigma error in the prediction of the Z boson mass in the Standard Model and a comparison of this result with experiment, leads to an error corridor, $\delta M_Z \sim 37$ MeV, where one can accommodate new physics. However, the more stringent constraint comes from fits to the high precision LEP data on the branching ratios of the Z decay and from the various asymmetries at the Z pole, when one demands that the χ^2 fits of StkSM are within 1% of that of the Standard Model. We will refer to this as the LEPI 1% constraint in the rest of the paper. Details of the method employed for the electroweak fits can be found in the analysis of Ref. [4, 5], and here we present just the results in the extended model with kinetic mixing.

Table 1 gives a fit to the LEP data for a specific point in the Stueckelberg parameter space with




FIG. 2: The plots give an analysis of the LEPII constraint. The upper plot, which has a Z' mass range of .2 – 1 TeV, shows the relevant LEPII contact interaction parameter $\Lambda_{VV}^{\pm l}$ as a function of χ^2 for the 19 observable of Table(1), where a $\chi^2 \sim 25$ is the SM fit as given in Table(1), and where the (yellow, red) shaded regions correspond to $\Lambda_{VV}^{\pm l} = (21.7, 17.1)$ TeV [30]. The lower plot for χ^2 in the range (25-50) gives $\Lambda_{VV}^{\pm l}$ as a function of the Z prime mass.

$\epsilon = .06$, $\delta = .03$, and $M_{Z'} \approx M_1 = 200$ GeV. In the analysis we have taken into account the constraint between g_Y and g_Y^{SM} and the inclusion of this constraint improves the electroweak fits over that of previous analyses for the case $\delta = 0$ [4, 5]. Thus the analysis of Table 1 shows that in the StkSM one finds χ^2 fits which are at the same level as in the SM. An analysis of χ^2 in the LEPI fits in the $\epsilon - \delta$ parameter space is given in Fig.(1). Specifically Fig.(1) shows that a large region of the parameter space can satisfy the LEPI 1% constraint. A striking aspect of the analysis of Fig.(1) is that this constraint is satisfied even though ϵ and δ can get significantly large, as long as $(\epsilon - \delta)$ is small. The physics of this is explained in Appendix A where it is shown that in the absence of matter in the hidden sector, there is only one effective parameter, $\bar{\epsilon}$, that enters the analysis of electroweak physics.

We discuss next the LEPII constraints. These constraints are typically characterized by the parameter of contact interaction Λ , and the LEPII group finds that $\Lambda_{VV} > (21.7, 17.1)$ TeV [30] to be the most constraining. The StkSM model predicts the theoretical value of Λ_{VV} through the following formula

$$\Lambda_{VV} = \frac{M_{Z'}}{M_Z} \sqrt{\frac{4\pi}{\sqrt{2}G_F v_e'^2}}. \quad (13)$$

TABLE I: Fits to 19 Z pole observables. Column 2 is given by the PDG [28], while the data in column 3 is from the SM Fit of the LEP EWWG [29]. The column labeled St Fit is an analysis for the input $\epsilon = 0.06$, $\delta = 0.03$, and $M_1 = 200$ GeV. In the last column PULL is defined by $(\text{Experiment} - \text{FIT})/\Delta$, and $\chi^2 = \sum \text{PULL}^2$.

Quantity	Experiment $\pm \Delta$	LEP FIT	St FIT	LEP PULL	St PULL
Γ_Z [GeV]	2.4952 ± 0.0023	2.4956	2.4956	-0.17	-0.17
σ_{had} [nb]	41.541 ± 0.037	41.476	41.469	1.76	1.95
R_e	20.804 ± 0.050	20.744	20.750	1.20	1.08
R_μ	20.785 ± 0.033	20.745	20.750	1.21	1.06
R_τ	20.764 ± 0.045	20.792	20.796	-0.62	-0.71
R_b	0.21643 ± 0.00072	0.21583	0.21576	0.83	0.93
R_c	0.1686 ± 0.0047	0.17225	0.17111	-0.78	-0.53
$A_{FB}^{(0,e)}$	0.0145 ± 0.0025	0.01627	0.01633	-0.71	-0.73
$A_{FB}^{(0,\mu)}$	0.0169 ± 0.0013	0.01627	0.01633	0.48	0.44
$A_{FB}^{(0,\tau)}$	0.0188 ± 0.0017	0.01627	0.01633	1.49	1.45
$A_{FB}^{(0,b)}$	0.0991 ± 0.0016	0.10324	0.10344	-2.59	-2.71
$A_{FB}^{(0,c)}$	0.0708 ± 0.0035	0.07378	0.07394	-0.85	-0.90
$A_{FB}^{(0,s)}$	0.098 ± 0.011	0.10335	0.10355	-0.49	-0.50
A_e	0.1515 ± 0.0019	0.1473	0.1476	2.21	2.05
A_μ	0.142 ± 0.015	0.1473	0.1476	-0.35	-0.37
A_τ	0.143 ± 0.004	0.1473	0.1476	-1.08	-1.15
A_b	0.923 ± 0.020	0.93462	0.93464	-0.58	-0.58
A_c	0.671 ± 0.027	0.66798	0.66812	0.11	0.11
A_s	0.895 ± 0.091	0.93569	0.93571	-0.45	-0.45
				$\chi^2 = 25.0$	$\chi^2 = 25.2$

A numerical analysis of the LEP II constraints is given in Fig.(1) and Fig.(2). The analysis of Fig.(1) exhibits that the LEPI 1% constraint is more stringent than the LEP II constraint, and thus the LEP II constraint is automatically satisfied once the LEPI 1% constraint is satisfied. This result is supported by the analysis of Fig.(2) which shows that the value of Λ_{VV} predicted by the model in the parameter space consistent with the LEPI 1% constraint is significantly larger than the lower limit of the LEP II constraint. The blue points that enter the shaded regions are eliminated by the LEP II constraint. However, these points also correspond to large χ^2 fits to the LEPI analysis and are eliminated by LEPI 1% constraint as well. Thus, for a narrow Z' , the LEPI 1% constraint is stronger than the LEP II constraint.

IV. MILLI-CHARGED DARK MATTER FROM THE HIDDEN SECTOR

In the previous section we did not include matter in the hidden sector which is defined as matter which is neutral under the SM gauge group but carries $U(1)_X$ quantum numbers and thus couples only to C_μ . The kinetic and mass mixings in the $U(1)_X \times U(1)_Y$ sectors typically generate milli-charges for such matter. The conditions for the origin of milli-charges arising from

such mixings are discussed in Appendix B, where simple examples are worked out to explain the constraints that lead to the appearance of such charges. Milli-charges have been examined in many works both theoretically and experimentally [7, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47]. Most of these analyses are in the context of kinetic mixing model of [7]. Here we consider the milli-charged matter in the hidden sector within the context of the Stueckelberg extension of the SM with both mass and kinetic mixing. If milli-charged matter exists then both the Z and the Z' can decay into it if kinematically allowed to do so. For the mass scales we investigate the milli-charge particle has a mass larger than $M_Z/2$. In this case all of the electroweak constraints discussed in Sec.III are unaffected. Further, the Z prime can decay into the millicharged matter if the mass of the hidden matter is less than $M_{Z'}/2$. Such decays increase the Z' width and thus decrease the branching ratios of the Z' decay into the visible sector which depletes the dilepton signal in the Drell-Yan process. A relatively strong dilepton signal manifests in the analysis of Refs. [3, 4, 5] where the Z' decays into the hidden sector were taken to be comparable to the Z' decays into the visible sector, i.e., $\Gamma_{Z'}^{\text{hid}} \sim \Gamma_{Z'}^{\text{vis}}$. This constraint then leads to a sharp Z' resonance, but the decay of the hidden sector matter via the Z' pole is not strong enough to annihilate the hidden sector milli-charged dark matter in




FIG. 3: An analysis of the relic density of milli-charged particles arising in the StSM Z' model from the hidden sector for the case $\delta = 0$, $M_\chi = 150$ GeV, $\epsilon = (.01 - .06)$ with .01 for the innermost curve and moving outward in steps of .01. The (yellow, grey) regions ($M_{Z'} < 2M_\chi$, $M_{Z'} > 2M_\chi$) correspond to a (narrow, broad) Z' resonance, and the WMAP-3 relic density constraints are satisfied for both a broad Z' resonance and a narrow Z' resonance as exhibited by the 2σ WMAP-3 red and black bands. The region of narrow Z' resonance is constrained by the LEP and Tevatron data. The region in the 2σ WMAP-3 band can be probed via a dilepton signal as shown in Fig.(4). The red band to the left is excluded by the CDF 95% C.L. [48] data while the black band is consistent or on the edge thereof, with all constraints which can produce an observable dilepton signal.

sufficient amounts to be consistent with experiment, unless extreme fine tuning is used (we return to this issue at the end of this section).

The recent work of Ref. [20] has carried out an explicit analysis of putting a pair of Dirac fermions in the hidden sector, and made the interesting observation that for values $g_X Q_X \leq O(1)$ the decay width of Z' into the hidden sector Dirac fermions (χ) can be of GeV size, and consequently the hidden matter can annihilate in sufficient amounts to satisfy the relic density. We have carried out a similar analysis using the thermal averaging procedure in the computation of the relic density as described in Appendix C. Our conclusions are in agreement with the analysis of Ref. [20] in the region of the parameter space investigated in Ref. [20] when no kinetic mixing is assumed in the absence of thermal averaging. In our work we take the kinetic mixing into account in the analysis of the relic density. We also make a further observation that there exists a significant region of the parameter space where it is possible to satisfy the relic density constraints and still have a narrow Z' resonance which can be detected at the Tevatron and at the




FIG. 4: An exhibition of the dilepton signal $\sigma \cdot Br(Z' \rightarrow l^+l^-)$ at the Tevatron consistent with the WMAP-3 relic density constraint as a function of $M_{Z'}$ when $2M_\chi = 300$ GeV. The curves in ascending order are for values of ϵ in the range $(0.01 - 0.06)$ in steps of 0.01. The dilepton signal has a dramatic fall as $M_{Z'}$ crosses the point $2M_\chi = 300$ GeV where the Z' decay into the hidden sector fermions is kinematically allowed, widening enormously the Z' decay width. The green shaded regions are where the WMAP-3 relic density constraints are satisfied for the case when there is no kinetic mixing. Red and blue regions are for the case when kinetic mixing is included. The DØ data set [49] was collected in the search for narrow resonances (RS) and is a stronger constraint to apply on this model than the recent CDF [48] data which put constraints on the parameter space when the Z' can decay only into matter in the visible sector.

LHC using the dilepton signal via a Drell-Yan process.

We give now further details of our relic density analysis. In the numerical analysis we will use $g_X = g_Y^{SM}$ and $Q_X = 1$ unless stated otherwise. We begin by considering the case when $\delta = 0$ which is the StSM case. Fig.(3) gives an analysis of the relic density as a function of $M_{Z'}$ for the case $M_\chi = 150$ GeV, and ϵ in the range $(0.01 - 0.06)$. Here one finds that the relic density is satisfied on two branches, one for $M_{Z'} > 2M_\chi$, and the other for $M_{Z'} < 2M_\chi$. In the region $M_{Z'} > 2M_\chi$ the relic density is satisfied over a broad range of Z' masses for appropriate ϵ values. In this region the decay width of the Z' is large, and thus the branching ratio into the visible sector is suppressed, which would make the dilepton signal from the Z' decay difficult to observe. This is exhibited in Fig.(4) where one finds that the dilepton signal essentially disappears in the region $M_{Z'} > 2M_\chi$. Next we examine the region $M_{Z'} < 2M_\chi$. Here one finds once again a satisfaction of the relic density even though the Z' mass lies below $2M_\chi$. The very sizable range of the Z' mass over which the relic density can be satisfied arises due to thermal averaging over the Z'

pole. As expected in this region the Z' will appear as a sharp narrow resonance since the decay of the Z' into the hidden sector fermions is kinematically disallowed. An analysis of the dilepton signal for this case is also given in Fig.(4), and a more detailed view is given in Fig.(5), which shows that the Drell-Yan signal $p\bar{p} \rightarrow Z' \rightarrow e^+e^-$ is enormously enhanced for $M_{Z'} < 2M_\chi$. Thus we have a region here of the parameter space where one will have a sharp resonance giving a visible dilepton signal while at the same time producing milli-charge dark matter consistent with WMAP-3.

We carry out a similar analysis for the case when the kinetic mixing parameter δ is non-vanishing. Fig.(6) gives an analysis of the effect of δ on the thermally averaged cross section $J(x_f)$, defined in Appendix C, as a function of the Z' mass. One finds that $J(x_f)$ is affected in a significant way by the change in δ and the particulars of the modification due to δ are explained in Appendix D. In Fig.(7) an analysis of relic density as a function of $M_{Z'}$ is given for $\bar{\epsilon} = 0.04$ and δ in the range $\delta = (.05 - .25)$. The analysis shows the relic density exhibits a strong δ dependence. Quite significantly, the region in $M_{Z'}$ over which the relic density can fall in the WMAP-3 region is widened. Once again, in the region $M_{Z'} > 2M_\chi$ the dilepton signal will be too dilute to be observable as shown in Fig.(7). On the other hand, in the region $M_{Z'} < 2M_\chi$, the Z' is a sharp narrow resonance since the Z' decay into the hidden sector fermions is kinematically




FIG. 5: Same as Fig.(4) except that only the mass region with a detectable dilepton signal at the Tevatron is exhibited but additional δ values are included in the analysis. The plots show that the region allowed by WMAP-3 constraints moves to the right for positive δ and to the left for negative δ . This phenomenon is explained in Appendix D. The inclusion of kinetic mixing is seen to enlarge the parameter space where the relic density constraints are satisfied and where an observable dilepton signal at the Tevatron can occur.

disallowed. Thus the dilepton signal here is strong as can be seen in Fig.(4) and Fig.(5) and should be observable with sufficient luminosity.

An analysis for the case of the LHC in the Drell-Yan process $pp \rightarrow Z' \rightarrow e^+e^-$, is given in Fig.(8). Here it is shown that the dilepton signal in the region $M_{Z'} < 2M_\chi$ consistent with the WMAP-3 constraints would be discernible with $10fb^{-1}$ of integrated luminosity for $\bar{\epsilon} \gtrsim .04$. The dilepton signal has a dramatic fall as $M_{Z'}$ crosses the point $M_{Z'} = 2M_\chi$ beyond which the Z' decay into the hidden sector fermions is kinematically allowed, widening enormously the Z' decay width. The green shaded regions are where the WMAP-3 relic density constraints are satisfied for the case when there is no kinetic mixing. Red and blue regions are for the case when kinetic mixing is included. When computing the dilepton signal, a 20 GeV mass bin is used, and a 50% detector cut is assumed. The thick solid line corresponds to the LHC discovery limit with $10fb^{-1}$ of integrated luminosity expected in the first year run, and the criteria used for the discovery limit is $5\sqrt{N_{SM}}$ or 10 events, whichever is larger, where N_{SM} is the number of SM background events. The analysis shows that data in the first year running at the LHC should be able to constrain the parameter space down to $\bar{\epsilon} \sim 0.04$ for $M_\chi = 500$ GeV when $M_{Z'} < 1$ TeV.

An interesting issue concerns the question regarding how small $\bar{\epsilon}$ can be for WMAP-3 relic density constraints to be satisfied. Fig.(9) addresses this question where an analysis of the relic density is given when $\bar{\epsilon} = 10^{-4}$. One finds satisfaction of the relic




FIG. 6: Illustration of the variation with δ of the thermally averaged cross section for the annihilation of milli-charged particles when $M_\chi = 250$ GeV and $\bar{\epsilon} = .03$. The generic effect of δ is to modify the shape of $J(x_f)$ and the region where the relic density constraints can be satisfied is enlarged. Further details are given in appendix D.




FIG. 7: An analysis of the relic density of milli-charged particles for the case when kinetic mixing is included in the Stueckelberg Z' model. The analysis is done for $M_\chi = 150$ GeV, $\bar{\epsilon} = .04$, and $\delta = (.05, .075, .10, .15, .20, .25)$, where the values are in descending order for $M_{Z'} > 300$ GeV. The red and black bands are the WMAP-3 constraints where the black band also produces an observable dilepton signal. The analysis shows that for $\bar{\epsilon}$ fixed, increasing δ increases the parameter space where the WMAP-3 relic density constraint is satisfied, while allowing for a detectable Z prime signal as shown in Figs.(4,5).

density in this case, and even smaller $\bar{\epsilon}$ were found admissible. Further, while the dilepton signal at the Tevatron in this case will be suppressed, it could still be visible at the LHC with sufficient luminosity.

Finally, we note that within the context of the Stueckelberg model it is possible to place indirect limits on the milli-charge coupling of hidden sector fermion with the photon from the Tevatron data. An analysis of the limits on the Stueckelberg mixing parameter ϵ was presented in [4] for the case $\delta = 0$. For this case, the milli-charge Q_{milli} , where $Q_{milli}e$ is the coupling of the photon with the hidden sector fermions (see Appendix D) is determined to be: $Q_{milli} \approx \epsilon$. Thus one may directly translate the limits obtained in [4] to limits on the milli-charged coupling of the hidden fermion with the photon. In the context of the present analysis, the cross-section predictions given here, (as for example, in Fig.(5)) with their overlapping WMAP-3 bands, shows this explicitly.

V. CONCLUSION

In the above we have given an analysis of the Stueckelberg extension of the Standard Model with




FIG. 8: An exhibition of the dilepton signal as given by the number of events for $10fb^{-1}$ of integrated luminosity at the LHC consistent with the WMAP-3 relic density constraint as a function of $M_{Z'}$ when $M_\chi = 500$ GeV. The curves in ascending order are for values of $\bar{\epsilon}$ in the range $(0.01 - 0.06)$ in steps of 0.01. Criteria set for the discovery limit are discussed in the text.




FIG. 9: An analysis of the relic density of milli-charged particles for the case of a very small $\bar{\epsilon}$. One finds that even for this case one can satisfy the WMAP-3 relic density constraints. In this case the hidden sector fermion mass is predicted to be essentially $M_{Z'}/2$. While the dilepton signal from the Drell-Yan process is too weak to be observable at the Tevatron, it is strong enough to be detectable at the LHC with sufficient integrated luminosity [5].

inclusion of the kinetic energy mixing in the $U(1)_X \times U(1)_Y$ sectors. Such kinetic mixings are quite generic in models with more than one $U(1)$ gauge group. It is shown that in the model with both the mass and kinetic mixing and in the absence of matter in the hidden sector, the sensitive parameter which measures the deviation from the Standard Model is given by $\bar{\epsilon}$

as defined by Eq. (22) which is a specific combination of ϵ and δ , where ϵ measures the mass mixing and δ measures the kinetic mixing. However, when matter in the hidden sector is taken into account, electroweak physics depends on both ϵ and δ . An analysis of the relic density of milli-charged dark matter which is generic in Stueckelberg extensions is given. Here our analysis is in agreement with the work of Ref. [20] for the case when no kinetic mixing is taken into account. Inclusion of the kinetic mixing is seen to increase the region of the parameter space where the relic density constraints consistent with WMAP-3 can be satisfied. We also analyze the Z' signal. As noted in Ref. [20] on the branch $M_{Z'} > 2M_\chi$ the dilepton signal from the Z' decay is too small to be observed at colliders, and our results are in agreement with this analysis. However, we note that on the branch $M_{Z'} < 2M_\chi$, there is a significant region of the parameter space where

the relic density constraints can be satisfied and the dilepton signal from the Z' decay via the Drell-Yan process is strong enough to be observed at the Tevatron and at the LHC. The analysis also shows that relic density constraints can be satisfied for values of $\bar{\epsilon}$ as low as 10^{-4} and even smaller values are possible. An interesting issue concerns the detectability of such dark matter in laboratory experiments which might put further limits on the parameter space or on the component of the relic density such matter can constitute. However, such an investigation is outside the scope of this work.

Acknowledgments

This work was supported in part by the U.S. National Science Foundation under the grant NSF-PHY-0546568.

Appendix A: Details of the Stueckelberg extension with kinetic mixing

In this Appendix we give further details of the Stueckelberg extension with kinetic mixing. The mass matrix in the vector boson sector, after applying the transformation that diagonalizes the kinetic energy, is given by

$$M^2 = K^T M_{St}^2 K, \quad (14)$$

and we display its explicit form below

$$M^2 = \begin{pmatrix} M_1^2 \frac{(1-\epsilon\delta)^2}{1-\delta^2} + \frac{1}{4}\gamma^2 v^2 \frac{\delta^2(1-2\epsilon\delta+\epsilon^2)}{1-\delta^2} & M_1^2 \frac{\epsilon(1-\epsilon\delta)}{\sqrt{1-\delta^2}} - \frac{1}{4}\gamma^2 v^2 \frac{\delta(1-2\epsilon\delta+\epsilon^2)}{\sqrt{1-\delta^2}} & \frac{1}{4}g_2\gamma v^2 \frac{\delta\sqrt{1-2\epsilon\delta+\epsilon^2}}{\sqrt{1-\delta^2}} \\ M_1^2 \frac{\epsilon(1-\epsilon\delta)}{\sqrt{1-\delta^2}} - \frac{1}{4}\gamma^2 v^2 \frac{\delta(1-2\epsilon\delta+\epsilon^2)}{\sqrt{1-\delta^2}} & M_1^2 \epsilon^2 + \frac{1}{4}\gamma^2 v^2 (1-2\epsilon\delta+\epsilon^2) & -\frac{1}{4}g_2\gamma v^2 \sqrt{1-2\epsilon\delta+\epsilon^2} \\ \frac{1}{4}g_2\gamma v^2 \frac{\delta\sqrt{1-2\epsilon\delta+\epsilon^2}}{\sqrt{1-\delta^2}} & -\frac{1}{4}g_2\gamma v^2 \sqrt{1-2\epsilon\delta+\epsilon^2} & \frac{1}{4}g_2^2 v^2 \end{pmatrix}. \quad (15)$$

The eigenvalues of Eq. (15) are

$$M_\gamma^2 = 0, \quad M_Z^2 = (q-p)/2, \quad M_{Z'}^2 = (q+p)/2, \quad (16)$$

where

$$p = \sqrt{\left(M_1^2\beta + \frac{\gamma^2\beta + g_2^2 v^2}{4}\right)^2 - 4M_1^2 \frac{\gamma^2 + g_2^2}{4} v^2 \beta},$$

$$q = M_1^2\beta + \frac{\gamma^2\beta + g_2^2 v^2}{4} \quad (17)$$

and where $\beta = (1-2\epsilon\delta+\epsilon^2)/(1-\delta^2)$. As expected in the limit $\epsilon = 0 = \delta$ the Stueckelberg sector decouples from the SM sector and one gets

$$M_Z = \frac{\sqrt{g_2^2 + \gamma^2}}{2} v, \quad M_{Z'} = M_1. \quad (18)$$

The expression for M_Z of Eq. (18) is exactly the result in the SM. However, quite remarkably decoupling takes place also in the limit when ϵ and δ are non-vanishing but satisfy the relation $\epsilon = \delta$. Here, while

the ϵ, δ dependence persists in the M^2 mass matrix of Eq. (15), the eigenvalues are still given by Eq. (18) in this limit. Further, the coupling of the Z boson to quarks and leptons reduces to that of the SM in this limit. One may understand this result by examining the explicit form of the transformation matrix of Eq. (7) which for the case $\epsilon = \delta$ becomes

$$T = \begin{pmatrix} 1 & \frac{\gamma}{\sqrt{g_2^2 + \gamma^2}} \frac{\delta}{\sqrt{1-\delta^2}} & -\frac{g_2}{\sqrt{g_2^2 + \gamma^2}} \frac{\delta}{\sqrt{1-\delta^2}} \\ 0 & -\frac{\gamma}{\sqrt{g_2^2 + \gamma^2}} \frac{1}{\sqrt{1-\delta^2}} & \frac{g_2}{\sqrt{g_2^2 + \gamma^2}} \frac{1}{\sqrt{1-\delta^2}} \\ 0 & \frac{g_2}{\sqrt{g_2^2 + \gamma^2}} & \frac{\gamma}{\sqrt{g_2^2 + \gamma^2}} \end{pmatrix}. \quad (19)$$

A direct substitution in the Z and Z' couplings then shows that they are identical to the case of the SM. These analytic forms explain the limits seen numerically in the analysis of Fig.(1).

Next we consider the case with arbitrary mass mixing and kinetic mixing. Here before diagonalizing the

mass matrix it is convenient to perform the following orthogonal transformation (O) given by

$$O = \begin{pmatrix} \sqrt{1-\delta^2} & -\delta & 0 \\ \delta & \sqrt{1-\delta^2} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (20)$$

which transforms the mass matrix to $\mathcal{M}^2 = O^T M^2 O$

$$\mathcal{M}^2 = \begin{pmatrix} M_1^2 & M_1^2 \bar{\epsilon} & 0 \\ M_1^2 \bar{\epsilon} & M_1^2 \bar{\epsilon}^2 + \frac{v^2}{4} \gamma^2 (1 + \bar{\epsilon}^2) & -\frac{v^2}{4} g_2 \gamma \sqrt{1 + \bar{\epsilon}^2} \\ 0 & -\frac{v^2}{4} g_2 \gamma \sqrt{1 + \bar{\epsilon}^2} & \frac{v^2}{4} g_2^2 \end{pmatrix} \quad (21)$$

where $\bar{\epsilon}$ is defined so that

$$\bar{\epsilon} = \frac{\epsilon - \delta}{\sqrt{1 - \delta^2}}. \quad (22)$$

We note that the mass matrix Eq. (21) looks exactly the same as for the mass matrix one has if there was just the Stueckelberg mass mixing except that ϵ is replaced by $\bar{\epsilon}$. Thus the orthogonal matrix which diagonalizes Eq. (21) will be the same as for the StSM case except that the free parameter is changed from ϵ to $\bar{\epsilon}$. We parameterize the rotation matrix $\mathcal{R}$ defined by $\mathcal{R}^T \mathcal{M}^2 \mathcal{R} = \text{Diag}(M_{Z'}^2, M_Z^2, 0)$ by the following

$$\mathcal{R} = \begin{bmatrix} \cos \psi \cos \phi - \sin \theta \sin \phi \sin \psi & \sin \psi \cos \phi + \sin \theta \sin \phi \cos \psi & -\cos \theta \sin \phi \\ \cos \psi \sin \phi + \sin \theta \cos \phi \sin \psi & \sin \psi \sin \phi - \sin \theta \cos \phi \cos \psi & \cos \theta \cos \phi \\ -\cos \theta \sin \psi & \cos \theta \cos \psi & \sin \theta \end{bmatrix}, \quad (23)$$

where the angles are defined so that

$$\tan \theta = \frac{\gamma}{g_2}, \quad \tan \phi = \bar{\epsilon}, \quad \tan 2\psi = \frac{2 \sin \theta M_0^2 \bar{\epsilon}}{M_1^2 - M_0^2 + (M_1^2 + M_0^2 - M_W^2) \bar{\epsilon}^2}, \quad (24)$$

and $M_0 = M_Z(\epsilon = \delta) = v\sqrt{g_2^2 + \gamma^2}/2$, $M_W = g_2 v/2$. The transformation relating the initial basis and the final diagonal basis is $V = [K(\delta)O(\delta)\mathcal{R}(\bar{\epsilon})]E$, where $V^T = (C, B, A^3)$, and $E^T = (Z', Z, A_\gamma)$. The neutral current interaction can now be written in the form

$$\mathcal{L}_{NC} = J^T S(\bar{\epsilon}, \delta) \mathcal{R}(\bar{\epsilon}) E \quad (25)$$

where $J^T = (g_X J_X, \gamma J_Y, g_2 J_2^3)$, and S is given by

$$S(\bar{\epsilon}, \delta) = \begin{bmatrix} 1 & -\delta/\sqrt{1-\delta^2} & 0 \\ 0 & \sqrt{1+\bar{\epsilon}^2} & 0 \\ 0 & 0 & 1 \end{bmatrix}. \quad (26)$$

When $J_X = 0$, one finds that the neutral current interaction of Eq. (25) has no dependence on δ . Expressing the tree level interaction in terms of the reduced vector and axial vector couplings as defined in Eq. (11), we find

$$v_f = \cos \psi [(1 - \bar{\epsilon} \sin \theta \tan \psi) T_f^3 - 2 \sin^2 \theta (1 - \bar{\epsilon} \csc \theta \tan \psi) Q_f], \quad (27)$$

$$a_f = \cos \psi [1 - \bar{\epsilon} \sin \theta \tan \psi] T_f^3, \quad (28)$$

$$v'_f = -\cos \psi [(\tan \psi + \bar{\epsilon} \sin \theta) T_f^3 - 2 \sin^2 \theta (\bar{\epsilon} \csc \theta + \tan \psi) Q_f], \quad (29)$$

$$a'_f = -\cos \psi [\tan \psi + \bar{\epsilon} \sin \theta] T_f^3. \quad (30)$$

Thus in the absence of the hidden sector matter, the effect on the neutral current sector of including the kinetic mixing in the Stueckelberg model, is the replacement of ϵ by $\bar{\epsilon}$, and in the limit $\bar{\epsilon} \rightarrow 0$ the Z couplings of the SM are recovered. However, with the inclusion of matter in the hidden, i.e., $J_X \neq 0$, the neutral current sector will depend on both $\bar{\epsilon}$ and δ . We exhibit this below

$$\mathcal{L}_{NC} = \mathcal{L}_{NC}(J_X = 0) + g_X J_X^\mu \left[(\mathcal{R}_{11} - S_\delta \mathcal{R}_{21}) Z'_\mu + (\mathcal{R}_{12} - S_\delta \mathcal{R}_{22}) Z_\mu + (\mathcal{R}_{13} - S_\delta \mathcal{R}_{23}) A_\mu^\gamma \right] \quad (31)$$

where S_δ appears in Eq.(31), and is as defined in Sec.(II). In this case the decay modes of Z' depend

on both ϵ and δ and an analysis of the branching

ratios should reveal the presence of kinetic mixing.

Further, as discussed in Sec.(IV) and in Appendix D, the relic density analysis depends sensitively on δ .

Appendix B: On the origin of milli-charged matter

In this Appendix we illustrate the mechanism which generates the milli-charge in the context of the analysis of this paper. We start with the kinetic mixing model [7] with two gauge fields $A_{1\mu}, A_{2\mu}$ corresponding to the gauge groups $U(1)$ and $U(1)'$. We choose the following Lagrangian $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1$ where

$$\begin{aligned}\mathcal{L}_0 &= -\frac{1}{4}F_{1\mu\nu}F_1^{\mu\nu} - \frac{1}{4}F_{2\mu\nu}F_2^{\mu\nu} - \frac{\delta}{2}F_{1\mu\nu}F_2^{\mu\nu}, \\ \mathcal{L}_1 &= J'_\mu A_1^\mu + J_\mu A_2^\mu.\end{aligned}\quad (32)$$

To put the kinetic energy term in its canonical form, one may use the transformation

$$\begin{bmatrix} A_1^\mu \\ A_2^\mu \end{bmatrix} \rightarrow K_0 \begin{bmatrix} A'^\mu \\ A^\mu \end{bmatrix}, \quad K_0 = \begin{bmatrix} \frac{1}{\sqrt{1-\delta^2}} & 0 \\ \frac{-\delta}{\sqrt{1-\delta^2}} & 1 \end{bmatrix}. \quad (33)$$

However, the transformation that canonically diagonalizes the kinetic energy is not unique. Thus, for example, $K = K_0 R$ instead of K_0 would do as well where R is an orthogonal matrix

$$R = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}. \quad (34)$$

Here $\mathcal{L}_1$ is given by

$$\begin{aligned}\mathcal{L}_1 &= A'^\mu \left[\frac{\cos \theta}{\sqrt{1-\delta^2}} J'_\mu + \left(\sin \theta - \frac{\cos \theta \delta}{\sqrt{1-\delta^2}} \right) J_\mu \right] \\ &+ A^\mu \left[-\frac{\sin \theta}{\sqrt{1-\delta^2}} J'_\mu + \left(\cos \theta + \frac{\sin \theta \delta}{\sqrt{1-\delta^2}} \right) J_\mu \right].\end{aligned}\quad (35)$$

In this case we see that each of the massless states interacts with the sources J and J' . However, one may choose θ to get asymmetric solutions. For instance for the case $\theta = \arctan [\delta/\sqrt{1-\delta^2}]$ one has

$$\mathcal{L}_1 = A^\mu \left[\frac{1}{\sqrt{1-\delta^2}} J_\mu - \frac{\delta}{\sqrt{1-\delta^2}} J'_\mu \right] + A'^\mu J'_\mu. \quad (36)$$

In this case while A' interacts only with the source J' , A interacts with both J and J' , with the coupling to the source J' proportional to the kinetic mixing parameter δ . We identify A with the physical photon field, J with the physical source arising from quarks

and leptons, while A' is the orthogonal massless state, and J' is the source in the hidden sector. Here the coupling of the photon with the hidden sector is proportional to δ and thus the hidden sector is milli-charged if δ is small.

Next we consider a model with kinetic mixing where a Stueckelberg mechanism generates a mass term of the type considered in Eq. (1)

$$\mathcal{L}_{\text{Mass}} = -\frac{1}{2}M_1^2 A_{1\mu} A_1^\mu - \frac{1}{2}M_2^2 A_{2\mu} A_2^\mu - M_1 M_2 A_{1\mu} A_2^\mu. \quad (37)$$

In this case diagonalization of the mass matrix fixes θ so that

$$\theta = \arctan \left[\frac{\epsilon \sqrt{1-\delta^2}}{1-\delta\epsilon} \right], \quad (38)$$

and the interaction Lagrangian is given by

$$\begin{aligned}\mathcal{L}_1 &= \frac{1}{\sqrt{1-2\delta\epsilon+\epsilon^2}} \left(\frac{\epsilon-\delta}{\sqrt{1-\delta^2}} J_\mu + \frac{1-\delta\epsilon}{\sqrt{1-\delta^2}} J'_\mu \right) A_M^\mu \\ &+ \frac{1}{\sqrt{1-2\delta\epsilon+\epsilon^2}} (J_\mu - \epsilon J'_\mu) A_\gamma^\mu.\end{aligned}\quad (39)$$

Here for the case $\epsilon = 0$ one finds that the massless state, the photon A_γ^μ , no longer couples with the hidden sector, while the massive mode A_M^μ couples with both the visible sector via J and with the hidden sector via J' . We conclude, therefore, that in the absence of the Stueckelberg mass mixing, for the case when only one mode is massless, there are no milli-charged particles coupled to the photon field. Thus milli-charge couplings appear in this case only when the Stueckelberg mixing parameter ϵ is introduced. Thus for the case when only one mode is massless the kinetic mixing by itself does not allow milli-charges but the Stueckelberg mass mixing model does.

Appendix C: Analysis of relic density of milli-charged dark matter in the hidden sector

The analysis of relic density involves the integral of the thermally averaged cross section from the current temperature to the freeze out such that [50]

$$J(x_f) = \int_0^{x_f} \langle \sigma v \rangle dx. \quad (40)$$

Here σ is the annihilation cross section for the process $\chi \bar{\chi} \rightarrow f \bar{f}$ where f denotes a quark or a lepton, and v is the relative velocity of the annihilating Dirac fermions χ and $\bar{\chi}$. We use the notation, $x = T/M_\chi$ in units where the Boltzman constant is unity, and x_f is the value of x at the freeze out temperature. The thermally averaged cross section $\langle \sigma v \rangle$ is given by

$$\langle \sigma v \rangle = \frac{\int_0^\infty dv v^2 (\sigma v) e^{-(v^2/4x)}}{\int_0^\infty dv v^2 e^{-(v^2/4x)}}. \quad (41)$$

Using the analysis of Ref. [20] $\langle\sigma v\rangle$ is given by

$$\langle\sigma v\rangle \simeq f_1(s)(\xi_L^2 + \xi_R^2) + f_2(s)(\xi_L\xi_R) \quad (42)$$

where the functions f_1 and f_2 depend on the square of the CM energy s and are given by

$$\begin{aligned} f_1(s) &= C_f v \frac{\beta_f}{s\beta_\chi} [s^2(1 + \frac{1}{3}\beta_f^2\beta_\chi^2) + 4M_\chi^2(s - 2m_f^2)] \\ f_2(s) &= C_f v \frac{\beta_f}{s\beta_\chi} [8m_f^2(s + 2M_\chi^2)]. \end{aligned} \quad (43)$$

Here $C_f = N_f/(32\pi)$, $N_f = (1, 3)$ for $f =$ (lepton, quark), ξ_L and ξ_R are as defined in Ref. [20] and contain the photon, the Z and the Z' poles, the latter augmented with a Breit-Wigner distribution for

thermal averaging, as the largest contribution to $\langle\sigma v\rangle$ for $4M_\chi^2 \gg M_Z^2$ arises from integration over the Z' pole. In the vicinity of the Z' pole we can expand $\langle\sigma v\rangle$ as follows

$$\langle\sigma v\rangle \simeq C_1 + C_2 v^2 + \frac{C_3 M_\chi^{-4} + C_4 M_\chi^{-2} v^2}{(v^2 - \epsilon_R)^2 + \gamma_R^2} \quad (44)$$

where ϵ_R and γ_R are given by

$$\epsilon_R = (M_{Z'}^2 - 4M_\chi^2)/M_\chi^2, \quad \gamma_R = \Gamma_{Z'} M_{Z'}/M_\chi^2 \quad (45)$$

and where $C_i (i = 1 - 4)$ can be read off from the expansion of s in Eq. (42) and Eq. (43). Using the technique of Ref. [50] the integral $J(x_f)$ is given by

$$J(x_f) = x_f C_1 + 3x_f^2 C_2 + \frac{1}{2\sqrt{\pi}M_\chi^4} \left[\frac{C_3}{\epsilon_R} I_2 + M_\chi^2 C_4 (I_1 + I_2) \right] \quad (46)$$

where

$$I_1 = \frac{1}{2} \int_0^\infty dy y^{-1/2} e^{-y} \text{Ln} \left[\frac{(4x_f y - \epsilon_R)^2 + \gamma_R^2}{\epsilon_R^2 + \gamma_R^2} \right], \quad (47)$$

$$I_2 = \frac{\epsilon_R}{\gamma_R} \int_0^\infty dy y^{-1/2} e^{-y} \left[\arctan \left[\frac{4x_f y - \epsilon_R}{\gamma_R} \right] + \arctan \left[\frac{\epsilon_R}{\gamma_R} \right] \right]. \quad (48)$$

From the above one then obtains the relic density using the standard relations as discussed, for example, in Ref. [50].

Appendix D: The effect of δ on the cross section

The kinetic mixing parameter δ enters Eq. (31) via the hidden sector where

$$\mathcal{L}_{NC}^{\text{hid}} = \bar{\chi} \gamma^\mu [\epsilon_{Z'}^\chi Z'_\mu + \epsilon_Z^\chi Z_\mu + \epsilon_\gamma^\chi A_\mu] \chi. \quad (49)$$

In the limit $\bar{\epsilon}, \delta \ll 1$, the couplings to leading order are

$$\epsilon_{Z'}^\chi = g_X Q_X \left[\mathcal{R}_{11} - \frac{\delta}{\sqrt{1-\delta^2}} \mathcal{R}_{21} \right] \simeq g_X Q_X, \quad (50)$$

$$\epsilon_Z^\chi = g_X Q_X \left[\mathcal{R}_{12} - \frac{\delta}{\sqrt{1-\delta^2}} \mathcal{R}_{22} \right] \simeq \bar{\epsilon} g_X Q_X \sin \theta \left[1 + \frac{\delta}{\bar{\epsilon}} \right], \quad (51)$$

$$\epsilon_\gamma^\chi = g_X Q_X \left[\mathcal{R}_{13} - \frac{\delta}{\sqrt{1-\delta^2}} \mathcal{R}_{23} \right] \simeq -\bar{\epsilon} g_X Q_X \cos \theta \left[1 + \frac{\delta}{\bar{\epsilon}} \right]. \quad (52)$$

For the purpose of illustration in this Appendix, and only here, we will work in the limit $m_f \sim 0$, $v \sim 0$ and $\Gamma_{Z'} \sim 0$ so that

$$\sigma v \simeq \frac{M_\chi^2 N_f}{2\pi} \left[\xi_L^2 + \xi_R^2 \right], \quad (53)$$

where for $4M_\chi^2 \gg M_Z^2$,

$$\begin{aligned}\xi_{L,R} &\simeq \frac{\epsilon_\gamma^\chi e Q_{\text{em}}^f}{4M_\chi^2} + \frac{\epsilon_Z^\chi \epsilon_Z^{fL,R}}{4M_\chi^2 - M_Z^2} + \frac{\epsilon_{Z'}^\chi \epsilon_{Z'}^{fL,R}}{4M_\chi^2 - M_{Z'}^2} \\ &= \alpha_{L,R} + \frac{\beta_{L,R}}{4M_\chi^2 - M_{Z'}^2}.\end{aligned}\quad (54)$$

Defining $x = (4M_\chi^2 - M_{Z'}^2)$, σv can be further simplified as

$$\sigma v \simeq a + \frac{2b}{x} + \frac{c}{x^2}, \quad (55)$$

where a, b, c are functions of $\eta = (1 + \delta/\bar{\epsilon})$ and are given by

$$\begin{aligned}a &= \frac{M_\chi^2 N_f}{2\pi} (\alpha_L^2 + \alpha_R^2) \\ b &= \frac{M_\chi^2 N_f}{2\pi} (\alpha_L \beta_L + \alpha_R \beta_R), \\ c &= \frac{M_\chi^2 N_f}{2\pi} (\beta_L^2 + \beta_R^2).\end{aligned}\quad (56)$$

The η dependence can then be read off so that $a(\eta) = \eta^2 a(1)$, $b(\eta) = \eta b(1)$, $c(\eta) = c(1)$. The cross section described by Eq. (55) has a dip when $x_{\min} = -c/b$ which gives $(\sigma v)_{\min} = a - b^2/c$. Now $x_{\min}$ and $(\sigma v)_{\min}$ have an η dependence of the form $x_{\min}(\eta) = x_{\min}(1)/\eta$, and $(\sigma v)_{\min}(\eta) = \eta^2 (\sigma v)_{\min}(1)$ which implies that for the case $\eta > 1$ a larger η shrinks $x_{\min}$ and moves the dip closer to the resonance point. At the same time it also increases $(\sigma v)_{\min}$. More interestingly, a negative η will switch the dip to the other side of the resonance. These effects are illustrated in the analysis of Fig.(6). The analysis above explains why the inclusion of kinetic mixing enhances the region where the relic density constraint is satisfied.

-
- [1] B. K rs and P. Nath, Phys. Lett. B **586**, 366 (2004) [hep-ph/0402047].
 - [2] B. K rs and P. Nath, JHEP **0412** (2004) 005 [hep-ph/0406167].
 - [3] B. K rs and P. Nath, JHEP **0507**, 069 (2005) [hep-ph/0503208].
 - [4] D. Feldman, Z. Liu and P. Nath, Phys. Rev. Lett. **97**, 021801 (2006) [hep-ph/0603039].
 - [5] D. Feldman, Z. Liu and P. Nath, JHEP **0611**, 007 (2006) [arXiv:hep-ph/0606294].
 - [6] T. Appelquist, H. C. Cheng and B. A. Dobrescu, Phys. Rev. D **64**, 035002 (2001); M. Battaglia, A. Datta, A. De Roeck, K. Kong and K. T. Matchev, JHEP **0507**, 033 (2005) [arXiv:hep-ph/0502041]; G. Burdman, B. A. Dobrescu and E. Ponton, Phys. Rev. D **74**, 075008 (2006).
 - [7] B. Holdom, B. Holdom, Phys. Lett. B **166**, 196 (1986); Phys. Lett. B **259**, 329 (1991).
 - [8] J. Kumar and J. D. Wells, Phys. Rev. D **74**, 115017 (2006) [arXiv:hep-ph/0606183].
 - [9] W. F. Chang, J. N. Ng and J. M. S. Wu, Phys. Rev. D **74** (2006) 095005 [arXiv:hep-ph/0608068].
 - [10] A. Ferrogli , A. Lorca and J. J. van der Bij, arXiv:hep-ph/0611174.
 - [11] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
 - [12] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. Lett. **84**, 2080 (2000).
 - [13] D. M. Ghilencea, L. E. Ibanez, N. Irges and F. Quevedo, JHEP **0208** (2002) 016; D. M. Ghilencea, Nucl. Phys. B **648** (2003) 215.
 - [14] L. E. Ibanez, F. Marchesano and R. Rabadan, JHEP **0111** (2001) 002; R. Blumenhagen, V. Braun, B. K rs and D. L st, hep-th/0210083; I. Antoniadis, E. Kiritsis, J. Rizos and T. N. Tomaras, Nucl. Phys. B **660**, 81 (2003); C. Coriano, N. Irges and E. Kiritsis, arXiv:hep-ph/0510332; R. Blumenhagen, B. Kors, D. Lust and S. Stieberger, arXiv:hep-th/0610327.
 - [15] P. Anastasopoulos, M. Bianchi, E. Dudas and E. Kiritsis, JHEP **0611**, 057 (2006) [hep-th/0605225]. P. Anastasopoulos, T. P. T. Dijkstra, E. Kiritsis and A. N. Schellekens, Nucl. Phys. B **759**, 83 (2006) [hep-th/0605226].
 - [16] C. Coriano and N. Irges, arXiv:hep-ph/0612128; C. Coriano and N. Irges, arXiv:hep-ph/0612140; C. Coriano, N. Irges and S. Morelli, arXiv:hep-ph/0701010;
 - [17] P. Anastasopoulos, arXiv:hep-th/0701114.

- [18] D. Feldman, B. Kors and P. Nath, Phys. Rev. D **75**, 023503 (2007) [arXiv:hep-ph/0610133].
- [19] D. N. Spergel *et al.*, astro-ph/0603449.
- [20] K. Cheung and T. C. Yuan, JHEP **0703**, 120 (2007) [arXiv:hep-ph/0701107].
- [21] K. S. Babu, C. F. Kolda and J. March-Russell, Phys. Rev. D **57**, 6788 (1998), hep-ph/9710441; R. Foot and X. G. He, Phys. Lett. B **267**, 509 (1991).
- [22] K.R. Dienes, C. Kolda, and J. March-Russell, Nucl. Phys **492**, 104 (1997)
- [23] M. Cvetič and P. Langacker, Mod. Phys. Lett. A **11** (1996) 1247 [hep-ph/9602424]; A.E. Farragi and M. Masip, Phys. Lett. B **388** (1996) 524 [hep-ph/9604302].
- [24] V. Barger, C. Kao, P. Langacker and H. S. Lee, Phys. Lett. B **600**, 104 (2004) [hep-ph/0408120];
- [25] A. Leike, Phys. Rept. **317**, 143 (1999), hep-ph/9805494;
- [26] For early work on the Stueckelberg mechanism see, E.C.G. Stueckelberg, Helv. Phys. Acta. **11** (1938) 225; V. I. Ogievetskii and I.V. Polubarinov, JETP **14** (1962) 179; A. Chodos and F. Cooper, Phys. Rev. D **3**, 2461 (1971); M. Kalb and P. Ramond, Phys. Rev. D **9**, 2273 (1974); T. J. Allen, M. J. Bowick and A. Lahiri, Mod. Phys. Lett. A **6** (1991) 559.
- [27] P. Nath and M. Yamaguchi, Phys. Rev. D **60**, 116004 (1999) [arXiv:hep-ph/9902323].
- [28] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592**, 1 (2004).
- [29] [ALEPH Collaboration], hep-ex/0509008.
- [30] [LEP Collaboration], arXiv:hep-ex/0312023; [ALEPH Collaboration], arXiv:hep-ex/0609051.
- [31] H. Goldberg and L. J. Hall, Phys. Lett. B **174**, 151 (1986).
- [32] E. Golowich and R. W. Robinett, Phys. Rev. D **35**, 391 (1987).
- [33] R. N. Mohapatra and I. Z. Rothstein, Phys. Lett. B **247**, 593 (1990).
- [34] S. Davidson and M. E. Peskin, Phys. Rev. D **49**, 2114 (1994) [arXiv:hep-ph/9310288].
- [35] R. Foot, G. C. Joshi, H. Lew and R. R. Volkas, Mod. Phys. Lett. A **5**, 95 (1990) [Erratum-ibid. A **5**, 2085 (1990)].
- [36] D. O. Caldwell, R. M. Eisberg, D. M. Grumm, M. S. Witherell, B. Sadoulet, F. S. Goulding and A. R. Smith, Phys. Rev. Lett. **61**, 510 (1988).
- [37] M. I. Dobroliubov and A. Y. Ignatiev, Phys. Rev. Lett. **65**, 679 (1990).
- [38] R. Foot, arXiv:hep-ph/9407331.
- [39] S. Davidson, S. Hannestad and G. Raffelt, JHEP **0005**, 003 (2000) [arXiv:hep-ph/0001179].
- [40] M. L. Perl, P. C. Kim, V. Halyo, E. R. Lee, I. T. Lee, D. Loomba and K. S. Lackner, Int. J. Mod. Phys. A **16**, 2137 (2001) [arXiv:hep-ex/0102033].
- [41] A. A. Prinz *et al.*, Phys. Rev. Lett. **81**, 1175 (1998) [arXiv:hep-ex/9804008].
- [42] S. L. Dubovsky, D. S. Gorbunov and G. I. Rubtsov, JETP Lett. **79**, 1 (2004) [Pisma Zh. Eksp. Teor. Fiz. **79**, 3 (2004)] [arXiv:hep-ph/0311189].
- [43] E. Masso and J. Redondo, Phys. Rev. Lett. **97**, 151802 (2006) [arXiv:hep-ph/0606163].
- [44] S. A. Abel and B. W. Schofield, Nucl. Phys. B **685**, 150 (2004) [arXiv:hep-th/0311051].
- [45] S. A. Abel, J. Jaeckel, V. V. Khoze and A. Ringwald, arXiv:hep-ph/0608248.
- [46] M. Ahlers, H. Gies, J. Jaeckel and A. Ringwald, arXiv:hep-ph/0612098.
- [47] A. Badertscher *et al.*, Phys. Rev. D **75**, 032004 (2007); S. N. Gninenko, N. V. Krasnikov and A. Rubbia, Phys. Rev. D **75**, 075014 (2007)
- [48] CDF Public Note : CDF/ANAL/EXOTIC/8421
- [49] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **95**, 091801 (2005) [arXiv:hep-ex/0505018].
- [50] P. Nath and R. Arnowitt, Phys. Rev. Lett. **70** (1993) 3696 [arXiv:hep-ph/9302318].; See also, H. Baer and M. Brhlik, Phys. Rev. D **53** (1996) 597 [arXiv:hep-ph/9508321] ; V. D. Barger and C. Kao, Phys. Rev. D **57** (1998) 3131 [arXiv:hep-ph/9704403]. For early work see, K. Griest and D. Seckel, Phys. Rev. D **43** (1991) 3191; P. Gondolo and G. Gelmini, Nucl. Phys. B **360** (1991) 145.